arXiv:1604.07613v4 [math.NT] 20 Sep 2016

HYPERGEOMETRIC FUNCTIONS AND ALGEBRAIC CURVES $y^e = x^d + ax + b$

PRAMOD KUMAR KEWAT AND RAM KUMAR

ABSTRACT. Let q be a prime power and $\mathbb{F}_q$ be a finite field with q elements. Let e and d be positive integers. In this paper, for $d \geq 2$ and $q \equiv 1 (\mathrm{mod}\ ed(d-1))$, we calculate the number of points on an algebraic curve $E_{e,d} : y^e = x^d + ax + b$ over a finite field $\mathbb{F}_q$ in terms of ${}_dF_{d-1}$ Gaussian hypergeometric series with multiplicative characters of orders d and $e(d-1)$, and in terms of ${}_{d-1}F_{d-2}$ Gaussian hypergeometric series with multiplicative characters of orders $ed(d-1)$ and $e(d-1)$. This helps us to express the trace of Frobenius endomorphism of an algebraic curve $E_{e,d}$ over a finite field $\mathbb{F}_q$ in terms of above hypergeometric series. As applications, we obtain some transformations and special values of ${}_2F_1$ hypergeometric series.

1. INTRODUCTION

In the 19^{th} century Gauss introduced classical hypergeometric series. Since then, many mathematicians such as Kummer, Ramanujan, Beukers, Stiller and others studied classical hypergeometric series extensively and found many interesting connections between classical hypergeometric series and different mathematical objects. In 1980's, Greene [7] introduced hypergeometric functions (or Gaussian hypergeometric series) over finite fields analogous to classical hypergeometric series as finite character sums over a finite field. It is found that these functions satisfy many summation and transformation formulas analogous to classical hypergeometric series. In a series papers, many interesting relations have been established between special values of these hypergeometric functions and number of points of certain algebraic curves over finite fields (see, for example, [1, 2, 3, 4, 5, 6, 8, 10, 12]).

Fuselier [6] and Lennon [10] found formulas for the trace of Frobenius endomorphism of certain family of elliptic curves in terms of Gaussian hypergeometric series with character of order 12. In [4], Barman and Kalita found the number of solutions of the polynomial equation $x^d + ax + b = 0$ over a finite field $\mathbb{F}_q$ in terms of special values of Gaussian hypergeometric series with characters of orders d and $d-1$ under the condition that $q \equiv 1 (\mathrm{mod}\ d(d-1))$ and $d \geq 2$. The same authors, in [5], expressed the number of $\mathbb{F}_q$-points on a hyperelliptic curve in terms of special values of Gaussian hypergeometric series. Let e and d be positive integers and $E_{e,d}$: $y^e = x^d + ax + b$ be an algebraic curve over a finite field $\mathbb{F}_q$. Throughout this paper we assume that $a, b \neq 0$. Let $N_{e,d}$ denotes the number of points on the algebraic curve $E_{e,d}$ over $\mathbb{F}_q$ excluding a point at infinity and $a_q(E_{e,d})$ denotes the trace of Frobenius of the algebraic curve $E_{e,d}$. In this paper, for $d \geq 2$ and $q \equiv 1 (\mathrm{mod}\ ed(d-1))$, we express $N_{e,d}$ and $a_q(E_{e,d})$ in terms of ${}_dF_{d-1}$ and ${}_{d-1}F_{d-2}$ Gaussian hypergeometric series containing multiplicative characters of orders d, $e(d-1)$ and $ed(d-1)$. We deduce the results of Lennon [10] on trace of Frobenius of elliptic curves from the main results. As an example, we also give an explicit formula for the trace of Frobenius of $E_{3,4}$ in terms of ${}_4F_3$ Gaussian hypergeometric series. As applications, we obtain some transformations and special values of ${}_2F_1$ hypergeometric series.

2. PRELIMINARIES

Let $\mathbb{F}_q$ be a finite field with q elements, where $q = p^n, p$ is a prime number and n is a positive integer. Note that $\mathbb{F}_q^* = \mathbb{F}_q \backslash \{0\}$ is a cyclic multiplicative group of order $q-1$. A multiplicative character $\chi \colon \mathbb{F}_q^* \longrightarrow \mathbb{C}^*$ is a group homomorphism. Throughout, we reserve the notations ε and ϕ for the trivial and the quadratic characters, respectively. Thus, for $x \in \mathbb{F}_q^*$

1991 *Mathematics Subject Classification.* 11G20, 11T24.

Key words and phrases. Algebraic curves, Gaussian hypergeometric series, Frobenius endomorphisms, Characters of finite fields.

$$\phi(x) = \left(\frac{x}{q}\right) = \begin{cases} 1, \text{ if } x \text{ is a square of some element in } \mathbb{F}_q^*, \\ -1, \text{if } x \text{ is not a square of any element in } \mathbb{F}_q^* \end{cases} \tag{2.1}$$

is the Legendre symbol. The following theorem gives the structure of multiplicative characters on $\mathbb{F}_q^*$.

Theorem 2.1. [11] *Let g be a generator of the multiplicative group $\mathbb{F}_q^*$. For $j = 0, 1, 2, \cdots, q-2$, the functions*

$$\chi_j(g^k) = e^{\frac{2\pi ijk}{q-1}}, \text{ for } k = 0, 1, 2, \cdots, q-2,$$

define multiplicative characters on $\mathbb{F}_q^*$.

The set $\widehat{\mathbb{F}_q^*}$ of all multiplicative characters on $\mathbb{F}_q^*$ is a cyclic group under multiplication of characters. One extends the domain of a multiplicative character χ on $\mathbb{F}_q^*$ to $\mathbb{F}_q$ by defining $\chi(0) = 0$.
Define the additive character $\theta : \mathbb{F}_q \to \mathbb{C}^*$ by $\theta(\alpha) = \zeta^{tr(\alpha)}$, where $\zeta = e^{\frac{2\pi i}{p}}$ and $tr : \mathbb{F}_q \to \mathbb{F}_p$ is the trace map given by

$$tr(\alpha) = \alpha + \alpha^p + \alpha^{p^2} + \cdots + \alpha^{p^{n-1}}.$$

Throughout this paper, capital letters A,B,C,$\cdots$ and Greek letters χ,ψ,$\cdots$ we will let denote multiplicative characters. Let δ denote both the function on $\mathbb{F}_q$ and the function on $\widehat{\mathbb{F}_q^*}$

$$\delta(x) = \begin{cases} 1 & \text{if } x = 0; \\ 0 & \text{if } x \neq 0, \end{cases} \tag{2.2}$$

and

$$\delta(A) = \begin{cases} 1 & \text{if } A \text{ is trivial character}, \\ 0 & \text{otherwise.} \end{cases} \tag{2.3}$$

Define $\overline{\chi}$ by $\chi\overline{\chi} = \epsilon$. We write $\sum_x$ to denote the sum over all x in $\mathbb{F}_q$ and $\sum_\chi$ to denote the sum over all characters of $\mathbb{F}_q$.
We recall the definitions of the Jacobi sum

$$J(A, B) = \sum_x A(x)B(1-x) \tag{2.4}$$

and the Gauss sum

$$G(A) =: \sum_x A(x)\zeta^{tr(x)}. \tag{2.5}$$

Definition 2.2. [11] *For characters A and B of $\mathbb{F}_q$, the binomial coefficient $\binom{A}{B}$ is defined as*

$$\binom{A}{B} = \frac{B(-1)}{q} J(A, \overline{B}).$$

In terms of the binomial coefficients, $A(1+x)$ can be written as

$$A(1+x) = \delta(x) + \frac{q}{q-1} \sum_\chi \binom{A}{\chi} \chi(x). \tag{2.6}$$

Some useful properties of the binomial coefficients which follow easily from properties of the Jacobi sums are

$$\binom{A}{B} = \binom{A}{A\overline{B}}, \tag{2.7}$$

$$\binom{A}{B} = \binom{B\overline{A}}{B} B(-1) \tag{2.8}$$

and

$$\binom{A}{B} = \binom{\overline{B}}{\overline{A}} AB(-1). \tag{2.9}$$

Let T be a fixed generator of $\widehat{\mathbb{F}_q^*}$ and $G_m = G(T^m)$. The following lemma gives the beautiful properties of the Gauss sum.

Lemma 2.3. [6] *If T^m is not trivial for $m \in \mathbb{N}$, then we have*

$$G_m G_{-m} = qT^m(-1).$$

The following lemmas give the nice relationship between the Gauss sum and the Jacobi sum.

Lemma 2.4. [10] *If T^{m-n} is not trivial for $m, n \in \mathbb{N}$, then we have*

$$\begin{aligned} G_m G_{-n} =& q\binom{T^m}{T^n} G_{m-n} T^n(-1) \\ =& J(T^m, T^{-n}) G_{m-n}. \end{aligned}$$

Lemma 2.5. [11] *If $T^{k_1}, T^{k_2}, \ldots, T^{k_n}$ are nontrivial multiplicative characters of $\mathbb{F}_q$ and $T^{k_1+k_2+\cdots+k_n}$ is nontrivial, then*

$$J(T^{k_1}, T^{k_2}, \ldots, T^{k_n}) = \frac{G_{k_1} G_{k_2} \ldots G_{k_n}}{G_{k_1+k_2+\cdots+k_n}}.$$

Lemma 2.6. [6] *Let T be a fixed generator of $\widehat{\mathbb{F}_q^*}$ and $n \in \mathbb{N}$, then orthogonality relations for multiplicative characters are given by*

(1) $\sum_{x \in \mathbb{F}_q^*} T^n(x) = \begin{cases} q-1 & \text{if } T^n = \epsilon; \\ 0 & \text{if } T^n \neq \epsilon. \end{cases}$

(2) $\sum_{n=0}^{q-2} T^n(x) = \begin{cases} q-1 & \text{if } x = 1; \\ 0 & \text{if } x \neq 1. \end{cases}$

The following lemma gives the relation between additive characters and the Gauss sums.

Lemma 2.7. [6] *Let θ be an additive character and $\alpha \in \mathbb{F}_q^*$, we have*

$$\theta(\alpha) = \frac{1}{q-1} \sum_{m=0}^{q-2} G_m T^m(\alpha).$$

Theorem 2.8. *(Davenport-Hasse Relation)*[9] *Let $m \in \mathbb{Z}^+$ and $q \equiv 1 (\mathrm{mod}\ m)$. For multiplication characters $\chi, \psi \in \widehat{\mathbb{F}_q^*}$, we have*

$$\prod_{\chi^m=1} G(\chi\psi) = -G(\psi^m)\psi(m^{-m}) \prod_{\chi^m=1} G(\chi).$$

We have the following two special cases of the above theorem.

Corollary 2.9. [4] *Let d be a positive integer, $l \in \mathbb{Z}, q \equiv 1(\mathrm{mod}\ \mathrm{d})$ and $t \in \{1, -1\}$. If $d > 1$ is an odd integer, then*

$$G_l G_{l+t\frac{q-1}{d}} G_{l+t\frac{2(q-1)}{d}} \cdots G_{l+t\frac{(d-1)(q-1)}{d}} = q^{\frac{d-1}{2}} T^{\frac{(d-1)(d+1)(q-1)}{8d}} (-1)\, T^{-l}\left(d^d\right) G_{ld}.$$

If d is an even integer, then

$$G_l G_{l+t\frac{q-1}{d}} G_{l+t\frac{2(q-1)}{d}} \cdots G_{l+t\frac{(d-1)(q-1)}{d}} = q^{\frac{d-2}{2}} G_{\frac{q-1}{2}} T^{\frac{(d-2)(q-1)}{8}} (-1) T^{-l}\left(d^d\right) G_{ld}.$$

The following lemma gives the value of the Gauss sums at the trivial and the quadratic characters,

Lemma 2.10. [6] *We have*

(1) $G(\epsilon) = G_0 = -1$

(2) $G(\phi) = G_{\frac{q-1}{2}} = \begin{cases} \sqrt{q} & \text{if } q \equiv 1(\mathrm{mod}\ 4); \\ i\sqrt{q} & \text{if } q \equiv 3(\mathrm{mod}\ 4). \end{cases}$

Theorem 2.11. *Let θ be an additive character and $x, y, z \in \mathbb{F}_q$. Then we have*

$$\sum_{z\in\mathbb{F}_q} \theta(z(x-y)) = q\delta(x,y),$$

where $\delta(x,y) = \begin{cases} 1 & \text{if } x = y; \\ 0 & \text{if } x \neq y. \end{cases}$

Definition 2.12. [11] *If $A_0, A_1, \cdots, A_n$ and $B_1, B_2, \cdots, B_n$ are characters of $\mathbb{F}_q$ and $x \in \mathbb{F}_q$, then the Gaussian hypergeometric series ${}_{n+1}F_n$ over $\mathbb{F}_q$ is defined as*

$${}_{n+1}F_n\left(\begin{matrix} A_0, & A_1, & \cdots, A_n \\ & B_1, & \cdots, B_n \end{matrix} \Big| x\right) = \frac{q}{q-1}\sum_{\chi} \binom{A_0\chi}{\chi}\binom{A_1\chi}{B_1\chi}\cdots\binom{A_n\chi}{B_n\chi}\chi(x).$$

3. main theorems

In this section, we state and prove the main theorems of this papers.

Theorem 3.1. *Let e and d be positive integers and let $N_{e,d}$ denote the number of $\mathbb{F}_q$ points on $E_{e,d} : y^e = x^d + ax + b$. If $d \geq 2$ and is an even integer, and $q \equiv 1(\mathrm{mod}(ed(d-1)))$, then*

$$N_{e,d} = q + \sum_{i=1}^{e-1} T^{-\frac{i(q-1)}{e}}(b) + \frac{1}{T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)} \sum_{i=1}^{e-1} M_i T^{\frac{i(q-1)}{e}} \left(\frac{d-1}{b}\right) \times$$

$${}_dF_{d-1}\left(\begin{matrix} \phi, & \epsilon, & \chi, & \chi^2, \cdots, \chi^{\frac{d-2}{2}}, & \chi^{\frac{d+2}{2}}, \cdots, \chi^{d-1} \\ & \psi^{(\frac{d}{2}e-i)}, & \psi^{e-i}, & \psi^{2e-i}, \cdots, \psi^{\frac{d-2}{2}e-i}, & \psi^{\frac{d+2}{2}e-i}, \cdots, \psi^{(d-1)e-i} \end{matrix} \Big| \alpha\right)$$

where $\alpha = \frac{d}{a}(\frac{bd}{a(d-1)})^{d-1}$, χ and ψ are characters of orders d and $e(d-1)$ respectively and

$$M_i = G_{-\frac{i(q-1)}{e}} G_{-\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}} G_{\frac{(id-e)(q-1)}{ed(d-1)}} G_{\frac{(id-2e)(q-1)}{ed(d-1)}} \cdots G_{\frac{(id-(\frac{d}{2}-1)e)(q-1)}{ed(d-1)}} G_{\frac{id-\left(\frac{d}{2}+1\right)e(q-1)}{e(d-1)}}$$
$$\cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}, i = 1, 2, \cdots, (e-1).$$

Remark 3.2. *In the above theorem, by using Lemmas 2.4 and 2.5, we can simplify the expression of M_i. If $e = 2$ then $M_1 = q^{\frac{d}{2}} T^{\frac{q-1}{2}}(-1)$ and if $e \neq 2$ then*

$$M_i = q^2 \frac{G_{\frac{(2i-e)(q-1)}{2e(d-1)}}}{G_{\frac{(id-\frac{d}{2}e)(q-1)}{ed(d-1)}}} \begin{pmatrix} T^{\frac{(2i-e)(q-1)}{2e}} \\ T^{\frac{i(q-1)}{e}} \end{pmatrix} \begin{pmatrix} T^{\frac{q-1}{2}} \\ T^{\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}} \end{pmatrix} T^{\frac{(2i(d-2)+ed)(q-1)}{2e(d-1)}}(-1)$$
$$J\left(T^{\frac{(id-e)(q-1)}{ed(d-1)}}, T^{\frac{(id-2e)(q-1)}{ed(d-1)}}, \ldots, T^{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}\right), i = 1, 2, \cdots, (e-1).$$

Theorem 3.3. *Let e and d be positive integers and let $N_{e,d}$ denote the number of $\mathbb{F}_q$ points on $E_{e,d} : y^e = x^d + ax + b$. If $d \geq 2$ and is an odd integer, and $q \equiv 1(\mathrm{mod}(ed(d-1)))$, then*

$$N_{e,d} =$$
$$q + \sum_{i=1}^{e-1} T^{-\frac{i(q-1)}{e}}(b) - N_i \frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{q^{(d-3)} G_{\frac{q-1}{2}}} \sum_{i=1}^{e-1} G_{-\frac{i(q-1)}{e}} T^{\frac{i(q-1)}{e}}(-1) T^{\frac{(e-i)(q-1)}{e}} \left(\frac{b}{d-1}\right)$$
$$+ \frac{T^{\frac{(4d^2+3d-1)(q-1)}{8d}}(-1)}{G_{\frac{q-1}{2}}} \sum_{i=1}^{e-1} G_{-\frac{i(q-1)}{e}} T^{\frac{-i(q-1)}{e}} \left(\frac{b}{d-1}\right) T^{\frac{(e-i)(q-1)}{e(d-1)}} \left(-\frac{1}{\alpha}\right) M_i \times$$

$${}_{d-1}F_{d-2}\left(\begin{matrix} \eta^{id-e}, & \eta^{de+id-2e}, \cdots, \eta^{\frac{ed^2-4ed+2id+e}{2}}, & \eta^{\frac{ed^2-2ed+2id-e}{2}}, \cdots, \eta^{ed^2-3ed+id+e} \\ & \psi^e, \cdots, \psi^{\frac{d-3}{2}e}, & \psi^{\frac{d-1}{2}e}, \cdots, \psi^{(d-2)e} \end{matrix} \Big| -\alpha\right)$$

where $\alpha = \frac{d}{a}(\frac{bd}{a(d-1)})^{d-1}$, η and ψ are characters of orders $ed(d-1)$ and $e(d-1)$ respectively,
$M_i = G_{\frac{(id-e)(q-1)}{ed(d-1)}} G_{\frac{(id-2e)(q-1)}{ed(d-1)}} \cdots G_{\frac{\left(id-\frac{d-1}{2}e\right)(q-1)}{ed(d-1)}} G_{\frac{\left(id-\frac{d+1}{2}e\right)(q-1)}{ed(d-1)}} \cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}$

and
$N_i = \{G_{\frac{q-1}{d}} G_{-\frac{(e-i)(q-1)}{e(d-1)}}\}\{G_{\frac{2(q-1)}{d}} G_{-\frac{(2e-i)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{2d}}\ G_{-\frac{(\frac{(d-1)}{2}e-i)(q-1)}{e(d-1)}}\}$
$\times \{G_{\frac{(d+1)(q-1)}{2d}} G_{-\frac{(\frac{d+1}{2}e-i)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{d}} G_{-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\},$
$i = 1, 2, \cdots, (e-1).$

Remark 3.4. *In the above theorem, by using Lemmas* 2.3, 2.4 *and* 2.5*, we can simplify the expressions of* M *and* N*. If* $e = 2$ *then* $M_1 = q^{\frac{d-1}{2}} T^{-\frac{(d-1)(q-1)}{8d}}(-1)$ *and* $N_1 = q^{d-1} T^{-\frac{(d-1)(q-1)}{8d}}(-1)$*. If* $e \neq 2$ *then*

$$M_i = G_{\frac{(2i-e)(q-1)}{2e}} J\left(T^{\frac{(id-e)(q-1)}{ed(d-1)}}, T^{\frac{(id-2e)(q-1)}{ed(d-1)}}, \ldots, T^{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}\right), i = 1, 2, \cdots, (e-1)$$

and

$$N_i = q^{\frac{d-1}{2}} T^{\frac{(d^2-1)(q-1)}{8d}}(-1) \quad G_{-\frac{(ed-2i)(q-1)}{2e}} J\left(T^{-\frac{(e-i)(q-1)}{e(d-1)}}, T^{-\frac{(2e-i)(q-1)}{e(d-1)}}, \ldots, T^{-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\right),$$
$$i = 1, 2, \cdots, (e-1).$$

Proof of the Theorem 3.1. Let $P(x, y) = x^d + ax + b - y^e$. Then $N_{e,d} = \#\{(x, y) \in \mathbb{F}_q^2 : P(x, y) = 0\}$. Now using the elementary identity,

$$\sum_{z \in \mathbb{F}_q} \theta(zP(x, y)) = \begin{cases} q & \text{if } P(x, y) = 0; \\ 0 & \text{if } P(x, y) \neq 0, \end{cases}$$

we obtain that

$$\begin{aligned} qN_{e,d} &= \sum_{x,y,z \in \mathbb{F}_q} \theta(zP(x, y)) \\ &= \sum_{x,y \in \mathbb{F}_q} \theta(0P(x, y)) + \sum_{z \in \mathbb{F}_q^*} \theta(zP(0, 0)) + \sum_{y,z \in \mathbb{F}_q^*} \theta(zP(0, y)) \\ &\quad + \sum_{x,z \in \mathbb{F}_q^*} \theta(zP(x, 0)) \sum_{x,y,z \in \mathbb{F}_q^*} \theta(zP(x, y)) \\ &= q^2 + \sum_{z \in \mathbb{F}_q^*} \theta(bz) + \sum_{y,z \in \mathbb{F}_q^*} \theta(bz)\theta(-zy^e) + \sum_{x,z \in \mathbb{F}_q^*} \theta(bz)\theta(zx^d)\theta(azx) \\ &\quad + \sum_{x,y,z \in \mathbb{F}_q^*} \theta(bz)\theta(zx^d)\theta(azx)\theta(-zy^e). \end{aligned}$$

This we can write as

$$qN_{e,d} := q^2 + A + B + C + D, \tag{3.1}$$

where

$$\begin{aligned} A &= \sum_{z \in \mathbb{F}_q^*} \theta(bz), \\ B &= \sum_{y,z \in \mathbb{F}_q^*} \theta(bz)\theta(-zy^e), \\ C &= \sum_{x,z \in \mathbb{F}_q^*} \theta(bz)\theta(zx^d)\theta(azx) \text{ and} \\ D &= \sum_{x,y,z \in \mathbb{F}_q^*} \theta(bz)\theta(zx^d)\theta(azx)\theta(-zy^e). \end{aligned}$$

By Lemma 2.7, A becomes

$$\begin{aligned}A =&\frac{1}{q-1}\sum_{l=0}^{q-2}G_{-l}\sum_{z\in\mathbb{F}_q^*}T^l(bz)\\ =&\frac{1}{q-1}\sum_{l=0}^{q-2}G_{-l}T^l(b)\sum_{z\in\mathbb{F}_q^*}T^l(z).\end{aligned}$$

This term is non zero only if $l=0$, using Part (1) of Lemmas 2.6 and 2.10, we have

$$A=-1.$$

Now again using Lemma 2.7, B becomes

$$B=\frac{1}{(q-1)^2}\sum_{l,m=0}^{q-2}G_{-l}G_{-m}T^l(b)T^m(-1)\sum_{z\in\mathbb{F}_q^*}T^{l+m}(z)\sum_{y\in\mathbb{F}_q^*}T^{em}(y).$$

This term is non zero only if $m=\frac{i(q-1)}{e}, i=0,1,\cdots(e-1)$ and $l=-m$. If $m=0$, then $l=0$ and if $m=\frac{i(q-1)}{e}, i=1,2,\cdots(e-1)$, then $l=-\frac{q-1}{e}$. By Part (1) of Lemmas 2.6 and 2.10, we have

$$\begin{aligned}B =&1+\sum_{i=1}^{e-1}G_{\frac{i(q-1)}{e}}G_{-\frac{i(q-1)}{e}}T^{-\frac{i(q-1)}{e}}(b)T^{\frac{i(q-1)}{e}}(-1)\\ =&1+q\sum_{i=1}^{e-1}T^{-\frac{i(q-1)}{e}}(b).\end{aligned}$$

Similarly, we calculate C and D,

$$C=\frac{1}{(q-1)^3}\sum_{l,m,n=0}^{q-2}G_{-l}G_{-m}G_{-n}T^l(b)T^n(a)\sum_{z\in\mathbb{F}_q^*}T^{l+m+n}(z)\sum_{x\in\mathbb{F}_q^*}T^{md+n}(x)$$

and

$$\begin{aligned}D =&\frac{1}{(q-1)^4}\sum_{l,m,n,k=0}^{q-2}G_{-l}G_{-m}G_{-n}G_{-k}T^l(b)T^n(a)T^k(-1)\sum_{z\in\mathbb{F}_q^*}T^{l+m+n+k}(z)\\ &\times\sum_{x\in\mathbb{F}_q^*}T^{md+n}(x)\sum_{y\in\mathbb{F}_q^*}T^{ek}(y).\end{aligned}$$

The innermost sum of D is non zero only if $k=\frac{i(q-1)}{e}, i=0,1,\cdots(e-1)$. For $k=0, D=-C$. Let $D_{\frac{i(q-1)}{e}}$ denote the value of D at $k=\frac{i(q-1)}{e}, i=1,2,\cdots(e-1)$. In this case

$$\begin{aligned}D_{\frac{i(q-1)}{e}} =&\frac{1}{(q-1)^3}\sum_{i=1}^{e-1}\sum_{l,m,n=0}^{q-2}G_{-l}G_{-m}G_{-n}G_{-\frac{i(q-1)}{e}}T^l(b)T^n(a)T^{\frac{i(q-1)}{e}}(-1)\\ &\times\sum_{z\in\mathbb{F}_q^*}T^{l+m+n+\frac{i(q-1)}{e}}(z)\sum_{x\in\mathbb{F}_q^*}T^{md+n}(x).\end{aligned}$$

Here, the term $D_{\frac{i(q-1)}{e}}=0$ unless $n=-md$ and $l=(d-1)(m+\frac{(e-i)(q-1)}{e(d-1)})$, thus,

$$\begin{aligned}D_{\frac{i(q-1)}{e}} =&\frac{1}{(q-1)}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)\sum_{m=0}^{q-2}G_{(d-1)\left(-m-\frac{(e-i)(q-1)}{e(d-1)}\right)}G_{-m}G_{md}\\ &\times T^{(d-1)\left(m+\frac{(e-i)(q-1)}{e(d-1)}\right)}(b)T^{-md}(a).\end{aligned}\tag{3.2}$$

By putting the values of A, B, C and D in Eq. (3.1), we get

$$qN_{e,d}=q^2+q\sum_{i=1}^{e-1}T^{-\frac{i(q-1)}{e}}(b)+D_{\frac{i(q-1)}{e}}.\tag{3.3}$$

If $d \geq 2$ is an even integer and $m \in \mathbb{Z}$, then Davenport-Hasse relation for $q \equiv 1 (\mathrm{mod}\ d),\ t \in \{1, -1\}$ gives

$$G_{dm} = \frac{G_m G_{m+\frac{q-1}{d}} G_{m+\frac{2(q-1)}{d}} \cdots G_{m+\frac{(d-1)(q-1)}{d}}}{q^{\frac{d-2}{2}} G_{\frac{q-1}{2}} T^{\frac{(d-2)(q-1)}{8}}(-1) T^{-m}(d^d)}$$

and

$$G_{(d-1)\left(-m-\frac{(e-i)(q-1)}{e(d-1)}\right)} = \frac{G_{-m-\frac{(e-i)(q-1)}{e(d-1)}} G_{-m-\frac{(2e-i)(q-1)}{e(d-1)}} \cdots G_{-m-\frac{((d-1)e-i)(q-1)}{e(d-1)}}}{q^{\frac{d-2}{2}} T^{\frac{d(d-2)(q-1)}{8(d-1)}}(-1) T^{m+\frac{(e-i)(q-1)}{e(d-1)}}(d-1)^{d-1}},$$

$i = 1, 2, \cdots, (e-1)$.

Now using G_{md} and $G_{(d-1)\left(-m-\frac{(e-i)(q-1)}{e(d-1)}\right)}$ in Eq.(3.2), we get

$$\begin{aligned} D_{\frac{i(q-1)}{e}} =& \frac{1}{q^{d-2}(q-1) G_{\frac{q-1}{2}} T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)} \sum_{i=1}^{e-1} \frac{G_{-\frac{i(q-1)}{e}} T^{\frac{(e-i)(q-1)}{e}}(b) T^{\frac{i(q-1)}{e}}(-1)}{T^{\frac{(e-i)(q-1)}{e}}(d-1)} \\ & \times \sum_{m=0}^{q-2} \{G_m G_{-m}\} \{G_{m+\frac{q-1}{d}} G_{-m-\frac{(e-i)(q-1)}{e(d-1)}}\} \{G_{m+\frac{2(q-1)}{d}} G_{-m-\frac{(2e-i)(q-1)}{e(d-1)}}\} \cdots \\ & \times \{G_{m+\frac{(d-2)(q-1)}{2d}} G_{-m-\frac{(\frac{d-2}{2}e-i)(q-1)}{e(d-1)}}\} \{G_{m+\frac{q-1}{2}} G_{-m-\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}\} \\ & \times \{G_{m+\frac{(d+2)(q-1)}{2d}} G_{-m-\frac{(\frac{d+2}{2}e-i)(q-1)}{e(d-1)}}\} \cdots \{G_{m+\frac{(d-1)(q-1)}{d}} G_{-m-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\} \\ & \times T^m \left(\frac{b^{d-1} d^d}{(d-1)^{d-1} a^d}\right). \end{aligned}$$

By arranging these terms, we get

$$\begin{aligned} D_{\frac{i(q-1)}{e}} =& \frac{1}{q^{d-2}(q-1) G_{\frac{q-1}{2}} T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)} \sum_{i=1}^{e-1} \frac{G_{-\frac{i(q-1)}{e}} T^{\frac{(e-i)(q-1)}{e}}(b) T^{\frac{i(q-1)}{e}}(-1)}{T^{\frac{(e-i)(q-1)}{e}}(d-1)} \\ & \times \sum_{m=0}^{q-2} \{G_{m+\frac{q-1}{2}} G_{-m}\} \{G_m G_{-m-\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}\} \{G_{m+\frac{q-1}{d}} G_{-m-\frac{(e-i)(q-1)}{e(d-1)}}\} \\ & \times \{G_{m+\frac{2(q-1)}{d}} G_{-m-\frac{(2e-i)(q-1)}{e(d-1)}}\} \cdots \{G_{m+\frac{(d-2)(q-1)}{2d}} G_{-m-\frac{(\frac{d-2}{2}e-i)(q-1)}{e(d-1)}}\} \\ & \times \{G_{m+\frac{(d+2)(q-1)}{2d}} G_{-m-\frac{(\frac{d+2}{2}e-i)(q-1)}{e(d-1)}}\} \cdots \{G_{m+\frac{(d-1)(q-1)}{d}} G_{-m-\frac{(d-1)e-i)(q-1)}{e(d-1)}}\} \\ & \times T^m \left(\frac{b^{d-1} d^d}{(d-1)^{d-1} a^d}\right). \end{aligned}$$

By using Lemma 2.4 in the above equation, we have

$$\begin{aligned}
D_{\frac{i(q-1)}{e}} =&\frac{q^2}{(q-1)G_{\frac{q-1}{2}}T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-\frac{d-1}{b})\sum_{m=0}^{q-2}G_{\frac{q-1}{2}}T^m(-1)\\
&\times\binom{T^{m+\frac{q-1}{2}}}{T^m}G_{-\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}T^{m+\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}(-1)\binom{T^m}{T^{m+\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}}G_{\frac{(id-e)(q-1)}{ed(d-1)}}\\
&\times T^{m+\frac{(e-i)(q-1)}{e(d-1)}}(-1)\binom{T^{m+\frac{q-1}{d}}}{T^{m+\frac{(e-i)(q-1)}{e(d-1)}}}G_{\frac{(id-2e)(q-1)}{ed(d-1)}}T^{m+\frac{(2e-i)(q-1)}{e(d-1)}}(-1)\binom{T^{m+\frac{2(q-1)}{d}}}{T^{m+\frac{(2e-i)(q-1)}{e(d-1)}}}\\
&\times\cdots G_{\frac{(id-(\frac{d}{2}-1)e)(q-1)}{ed(d-1)}}T^{m+\frac{(\frac{d-2}{2}e-i)(q-1)}{e(d-1)}}(-1)\binom{T^{m+\frac{(d-2)(q-1)}{2d}}}{T^{m+\frac{(\frac{id-2}{2}e-1)(q-1)}{e(d-1)}}}\\
&\times G_{\frac{(id-(\frac{d}{2}+1)e)(q-1)}{ed(d-1)}}T^{m+\frac{(\frac{d+2}{2}e-i)(q-1)}{e(d-1)}}(-1)\binom{T^{m+\frac{(d+2)(q-1)}{2d}}}{T^{m+\frac{(\frac{d+2}{2}e-i)(q-1)}{e(d-1)}}}\cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}\\
&\times T^{m+\frac{((d-1)e-i)(q-1)}{e(d-1)}}(-1)\binom{T^{m+\frac{(d-1)(q-1)}{d}}}{T^{m+\frac{((d-1)e-i)(q-1)}{e(d-1)}}}T^m\left(\frac{b^{(d-1)}d^d}{(d-1)^{(d-1)}a^d}\right).
\end{aligned}$$

By collecting the terms of G_s and T^s, we get

$$\begin{aligned}
D_{\frac{i(q-1)}{e}} =&\frac{q^2}{(q-1)T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)}\sum_{i=1}^{e-1}M_iT^{\frac{i(q-1)}{e}}(-\frac{d-1}{b})\sum_{m=0}^{q-2}T^{md+\frac{d(q-1)}{2}}(-1)\\
&\times T^{-\frac{i(q-1)}{e}}(-1)\binom{T^{m+\frac{q-1}{2}}}{T^m}\binom{T^m}{T^{m+\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}}\binom{T^{m+\frac{q-1}{d}}}{T^{m+\frac{(e-i)(q-1)}{e(d-1)}}}\binom{T^{m+\frac{2(q-1)}{d}}}{T^{m+\frac{(2e-i)(q-1)}{e(d-1)}}}\\
&\times\cdots\binom{T^{m+\frac{(d-2)(q-1)}{2d}}}{T^{m+\frac{\frac{d-2}{2}e-i)(q-1)}{e(d-1)}}}\binom{T^{m+\frac{(d+2)(q-1)}{2d}}}{T^{m+\frac{\frac{d+2}{2}e-i)(q-1)}{e(d-1)}}}\cdots\binom{T^{m+\frac{(d-1)(q-1)}{d}}}{T^{m+\frac{((d-1)e-i)(q-1)}{e(d-1)}}}T^m\left(\frac{b^{d-1}d^d}{(d-1)^{d-1}a^d}\right),
\end{aligned}$$

where

$$\begin{aligned}
M_i =&G_{-\frac{i(q-1)}{e}}G_{-\frac{(\frac{d}{2}e-i)(q-1)}{e(d-1)}}G_{\frac{(id-e)(q-1)}{ed(d-1)}}G_{\frac{(id-2e)(q-1)}{ed(d-1)}}\cdots G_{\frac{(id-(\frac{d}{2}-1)e)(q-1)}{ed(d-1)}}G_{\frac{id-\left(\frac{d}{2}+1\right)e(q-1)}{e(d-1)}}\\
&\cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}, i=1,2,\cdots,(e-1).
\end{aligned}$$

Since d is even, thus, $T^{md+\frac{d(q-1)}{2}}(-1)=1$. By Definition 2.12, we have

$$\begin{aligned}
D_{\frac{q-1}{e}} =&\frac{q}{T^{\frac{(d-2)(2d-1)(q-1)}{8(d-1)}}(-1)}\sum_{i=1}^{e-1}M_iT^{\frac{i(q-1)}{e}}\left(\frac{d-1}{b}\right)\\
\times_d F_{d-1}&\left(\begin{matrix}\phi, & \epsilon, & \chi, & \chi^2,\cdots,\chi^{\frac{d-2}{2}}, & \chi^{\frac{d+2}{2}},\cdots,\chi^{d-1}\\ & \psi^{(\frac{d}{2}e-i)}, & \psi^{e-i}, & \psi^{2e-i},\cdots,\psi^{\frac{d-2}{2}e-i}, & \psi^{\frac{d+2}{2}e-i},\cdots,\psi^{(d-1)e-i}\end{matrix}\Bigg|\alpha\right),
\end{aligned}$$

where $\alpha=\frac{d}{a}(\frac{bd}{a(d-1)})^{(d-1)}$, χ and ψ are characters of orders d and $e(d-1)$ respectively. We complete the proof by putting the value of $D_{\frac{i(q-1)}{e}}$ in equation (3.3). □

Proof of the Theorem 3.3. If $d\geq 2$ and is an odd integer, then Davenport-Hasse relations for G_{dm} and $G_{(d-1)(-m-\frac{(e-1)(q-1)}{e(d-1)})}$ are given by

$$G_{dm}=\frac{G_mG_{m+\frac{q-1}{d}}G_{m+\frac{2(q-1)}{d}}\cdots G_{m+\frac{(d-1)(q-1)}{d}}}{q^{\frac{d-1}{2}}T^{\frac{(d-1)(d+1)(q-1)}{8d}}(-1)T^{-m}(d^d)},$$

$$\begin{aligned}
G_{(d-1)(-m-\frac{(e-i)(q-1)}{e(d-1)})} =&\frac{G_{-m-\frac{(e-i)(q-1)}{e(d-1)}}G_{-m-\frac{(2e-i)(q-1)}{e(d-1)}}\cdots G_{-m-\frac{((d-1)e-i)(q-1)}{e(d-1)}}}{G_{\frac{q-1}{2}}q^{\frac{d-3}{2}}T^{\frac{(d-3)(q-1)}{8}}(-1)T^{m+\frac{(e-i)(q-1)}{e(d-1)}}(d-1)^{(d-1)}},\\
&i=1,2,\cdots,(e-1).
\end{aligned}$$

Using these identities in Eq. (3.2), we get

$$\begin{aligned}
D_{\frac{i(q-1)}{e}} =&\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{(q-1)q^{(d-2)}G_{\frac{q-1}{2}}T^{\frac{(d^2-1)(q-1)}{4d}}(-1)}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{(e-i)(q-1)}{e}}\left(\frac{b}{d-1}\right)\\
&\times\sum_{m=0}^{q-2}\{G_mG_{-m}\}\{G_{m+\frac{q-1}{d}}G_{-m-\frac{(e-i)(q-1)}{e(d-1)}}\}\{G_{m+\frac{2(q-1)}{d}}G_{-m-\frac{(2e-i)(q-1)}{e(d-1)}}\}\cdots\\
&\times\{G_{m+\frac{(d-1)(q-1)}{2d}}\ G_{-m-\frac{(\frac{(d-1)}{2}e-i)(q-1)}{e(d-1)}}\}\{G_{m+\frac{(d+1)(q-1)}{2d}}G_{-m-\frac{(\frac{d+1}{2}e-i)(q-1)}{e(d-1)}}\}\\
&\times\cdots\{G_{m+\frac{(d-1)(q-1)}{d}}G_{-m-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\}T^m\left(\frac{b^{(d-1)}d^d}{(d-1)^{(d-1)}a^d}\right).
\end{aligned}$$

Since d is an odd integer, thus $T^{\frac{(d^2-1)(q-1)}{4d}}(-1)=1$. Next, we eliminate the term $\{G_mG_{-m}\}$, using the fact that if $m=0$, then $G_mG_{-m}=qT^m(-1)-(q-1)$, and if $m\neq 0$, then $G_mG_{-m}=qT^m(-1)$. Using these identities in the above equation, we have

$$\begin{aligned}
D_{\frac{i(q-1)}{e}} =&\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{(q-1)q^{(d-2)}G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{(e-i)(q-1)}{e}}\left(\frac{b}{d-1}\right)\\
&\times\sum_{m=0}^{q-2}\{G_{m+\frac{q-1}{d}}G_{-m-\frac{(e-1)(q-1)}{e(d-1)}}\}\{G_{m+\frac{2(q-1)}{d}}G_{-m-\frac{(2e-1)(q-1)}{e(d-1)}}\}\cdots\\
&\{G_{m+\frac{(d-1)(q-1)}{2d}}\ G_{-m-\frac{\left(\frac{d-1}{2}e-1\right)(q-1)}{e(d-1)}}\}\{G_{m+\frac{(d+1)(q-1)}{2d}}G_{-m-\frac{\left(\frac{d+1}{2}e-1\right)(q-1)}{e(d-1)}}\}\\
&\times\cdots\{G_{m+\frac{(d-1)(q-1)}{d}}G_{-m-\frac{((d-1)e-1)(q-1)}{e(d-1)}}\}T^m\left(-\frac{b^{(d-1)}d^d}{(d-1)^{(d-1)}a^d}\right)\\
&-\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{q^{(d-2)}G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{(e-i)(q-1)}{e}}\left(\frac{b}{d-1}\right)\\
&\times\{G_{\frac{q-1}{d}}G_{-\frac{(e-i)(q-1)}{e(d-1)}}\}\{G_{\frac{2(q-1)}{d}}G_{-\frac{(2e-i)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{2d}}\ G_{-\frac{\left(\frac{d-1}{2}e-i\right)(q-1)}{e(d-1)}}\}\\
&\times\{G_{\frac{(d+1)(q-1)}{2d}}G_{-\frac{\left(\frac{d+1}{2}e-i\right)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{d}}G_{-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\}.
\end{aligned}$$

Using Lemma 2.4 in each terms of the above equation and collecting the terms of G_s and T^s. Since d is an odd integer, thus $T^{m(d-1)}(-1)=1$, we have

$$\begin{aligned}
D_{\frac{i(q-1)}{e}} =&\frac{q^2T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{(q-1)G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{-\frac{i(q-1)}{e}}(-1)T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)\\
&\times\{G_{\frac{(id-e)(q-1)}{ed(d-1)}}G_{\frac{(id-2e)(q-1)}{ed(d-1)}}G_{\frac{(id-3e)(q-1)}{ed(d-1)}}\cdots G_{\frac{\left(id-\frac{(d-1)}{2}e\right)(q-1)}{ed(d-1)}}G_{\frac{\left(id-\frac{(d+1)}{2}e\right)(q-1)}{ed(d-1)}}\\
&\times\cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}\}\sum_{m=0}^{q-2}\binom{T^{m+\frac{q-1}{d}}}{T^{m+\frac{(e-i)(q-1)}{e(d-1)}}}\binom{T^{m+\frac{2(q-1)}{d}}}{T^{m+\frac{(2e-i)(q-1)}{e(d-1)}}}\cdots\\
&\times\binom{T^{m+\frac{(d-1)(q-1)}{2d}}}{T^{m+\frac{\left(\frac{d-1}{2}e-i\right)(q-1)}{e(d-1)}}}\binom{T^{m+\frac{(d+1)(q-1)}{2d}}}{T^{m+\frac{\left(\frac{d+1}{2}e-i\right)(q-1)}{e(d-1)}}}\cdots\binom{T^{m+\frac{(d-1)(q-1)}{d}}}{T^{m+\frac{((d-1)e-i)(q-1)}{e(d-1)}}}T^m(-\alpha)\\
&-\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{q^{(d-2)}G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)N_i,
\end{aligned}$$

where

$$\begin{aligned}
N_i =&\{G_{\frac{q-1}{d}}G_{-\frac{(e-i)(q-1)}{e(d-1)}}\}\{G_{\frac{2(q-1)}{d}}G_{-\frac{(2e-i)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{2d}}\ G_{-\frac{\left(\frac{d-1}{2}e-i\right)(q-1)}{e(d-1)}}\}\\
&\times\{G_{\frac{(d+1)(q-1)}{2d}}G_{-\frac{\left(\frac{d+1}{2}e-i\right)(q-1)}{e(d-1)}}\}\cdots\{G_{\frac{(d-1)(q-1)}{d}}G_{-\frac{((d-1)e-i)(q-1)}{e(d-1)}}\},\\
&i=1,2,\cdots,(e-1),
\end{aligned}$$

and $\alpha=\left(\frac{d}{a}\right)\left(\frac{bd}{a(d-1)}\right)^{(d-1)}$. Replace $m+\frac{(e-i)(q-1)}{e(d-1)}$ by m in the above equation, we have

$$\begin{aligned}D_{\frac{i(q-1)}{e}}=&q^2\frac{T^{\frac{(4d^2+3d-1)(q-1)}{8d}}(-1)}{(q-1)G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)M_i\sum_{m=0}^{q-2}\binom{T^{m+\frac{(id-e)(q-1)}{ed(d-1)}}}{T^m}\\&\times\binom{T^{m+\frac{(id+(d-2)e)(q-1)}{ed(d-1)}}}{T^{m+\frac{e(q-1)}{e(d-1)}}}\cdots\binom{T^{m+\frac{(ed^2-4ed+2id+e)(q-1)}{2ed(d-1)}}}{T^{m+\frac{(d-3)e(q-1)}{2e(d-1)}}}\binom{T^{m+\frac{(ed^2-2ed+2id-e)(q-1)}{2ed(d-1)}}}{T^{m+\frac{(d-1)e(q-1)}{2e(d-1)}}}\\&\times\cdots\binom{T^{m+\frac{(ed^2-3ed+id+e)(q-1)}{ed(d-1)}}}{T^{m+\frac{(d-2)e(q-1)}{e(d-1)}}}T^m(-\alpha)T^{-\frac{(e-i)(q-1)}{e(d-1)}}(-\alpha)\\&-\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{q^{(d-2)}G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)N_i,\end{aligned}$$

where $M_i=\{G_{\frac{(id-e)(q-1)}{ed(d-1)}}G_{\frac{(id-2e)(q-1)}{ed(d-1)}}G_{\frac{(id-3e)(q-1)}{ed(d-1)}}\cdots G_{\frac{\left(id-\frac{d-1}{2}e\right)(q-1)}{ed(d-1)}}G_{\frac{\left(id-\frac{d+1}{2}e\right)(q-1)}{ed(d-1)}}$
$\times G_{\frac{\left(id-\frac{d+3}{2}e\right)(q-1)}{ed(d-1)}}\cdots G_{\frac{(id-(d-1)e)(q-1)}{ed(d-1)}}\},i=1,2,\cdots,(e-1)$.
using Definition 2.12, in the above equation, we have

$$\begin{aligned}D_{\frac{i(q-1)}{e}}=&-N_i\frac{T^{\frac{(3d-1)(q-1)}{8d}}(-1)}{q^{(d-2)}G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{i(q-1)}{e}}(-1)T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)\\&+q\frac{T^{\frac{(4d^2+3d-1)(q-1)}{8d}}(-1)}{G_{\frac{q-1}{2}}}\sum_{i=1}^{e-1}G_{-\frac{i(q-1)}{e}}T^{\frac{-i(q-1)}{e}}\left(\frac{b}{d-1}\right)M_iT^{-\frac{(e-i)(q-1)}{e(d-1)}}(-\alpha)\end{aligned}$$

$$\times{}_{d-1}F_{d-2}\left(\begin{matrix}\eta^{id-e}, & \eta^{id+ed-2e},\cdots,\eta^{\frac{ed^2-4ed+2id+e}{2}}, & \eta^{\frac{ed^2-2ed+2id-e}{2}},\cdots,\eta^{ed^2-3ed+id+e}\\ & \psi^e,\cdots,\psi^{\frac{d-3}{2}e}, & \psi^{\frac{d-1}{2}e},\cdots,\psi^{(d-2)e}\end{matrix}\Bigg|-\alpha\right),$$

where η, ψ are characters of orders $ed(d-1)$ and $e(d-1)$ respectively. We complete the proof by putting the value of $D_{\frac{q-1}{e}}$ in equation (3.3). Note that if $e=2$, then N_1 becomes $q^{d-1}T^{-\frac{(d-1)(q-1)}{8d}}(-1)$ and $M_1=q^{\frac{d-1}{2}}T^{-\frac{(d-1)(q-1)}{8d}}(-1)$. □

Remark 3.5. *Let q,e,d and $E_{e,d}$ be as in Theorems* 3.1 *and* 3.3. *Let $a_q(E_{e,d}(\mathbb{F}_q))$ denotes the trace of Frobenius of the algebraic curve $E_{e,d}$. Since $a_q(E_{e,d}(\mathbb{F}_q))=q-N_{e,d}$, from Theorems* 3.1 *and* 3.3, *we can express the trace of Frobenius of the algebraic curve $E_{e,d}$ in terms of ${}_dF_{d-1}$ and ${}_{d-1}F_{d-2}$ Gaussian hypergeometric series containing multiplicative characters of orders d, $e(d-1)$ and $ed(d-1)$.*

4. Applications

In the following example, we deduce Theorem 2.1 of Lennon [10] from Theorem 3.3.

Example 4.1. [10] *Let $q=p^n,p>3$ a prime and $q\equiv1(\mathrm{mod}\ 12)$, Let $E_{2,3}:y^2=x^3+ax+b$ be an elliptic curve over $\mathbb{F}_q$ with $j(E_{2,3})\neq0,1728$, then the trace of Frobenius map on $E_{2,3}$ can be expressed as*

$$a_q(E_{2,3})=-qT^{\frac{q-1}{4}}\left(\frac{a^3}{27}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & T^{\frac{q-1}{2}}\end{matrix}\Bigg|-\frac{27b^2}{4a^3}\right).$$

Let $e=2$ and $d=3$ in Theorem 3.3. *In this case, we have $M_1=qT^{-\frac{q-1}{12}}(-1)$, $N_1=q^2T^{-\frac{q-1}{12}}(-1)$ and*
$N_{2,3}=q+\phi(b)-T^{-\frac{q-1}{12}}(-1)\phi(-2b)T^{\frac{q-1}{3}}(-1)+\frac{\phi(-2b)T^{\frac{q-1}{3}}(-1)qT^{\frac{-q-1}{3}}(-1)}{T^{\frac{q-1}{4}}\left(-\frac{27b^2}{4a^3}\right)}$

$$\times {}_2F_1\left(\begin{matrix}\eta, & \eta^5 \\ & \psi^2\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right),$$

where η is a multiplicative character of order 12 and ψ is multiplicative character of order 4. Thus, we have

$$\begin{aligned}N_{2,3} =& q+\phi(b)-\phi(-2b)T^{\frac{q-1}{4}}(-1)+\frac{qT^{\frac{q-1}{4}}(-1)}{T^{\frac{q-1}{4}}\left(-\frac{27}{a^3}\right)}{}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right)\\ =& q+\phi(b)-\phi(-2b)T^{\frac{q-1}{4}}(-1)+qT^{\frac{q-1}{4}}(-1)T^{\frac{q-1}{4}}\left(-\frac{a^3}{27}\right)\\ & \times {}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right)\\ =& q+\phi(b)-\phi(-2b)T^{\frac{q-1}{4}}(-1)+qT^{\frac{q-1}{4}}\left(\frac{a^3}{27}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right).\end{aligned}$$

Since $\phi(-2b)=T^{\frac{q-1}{2}}(-2)\phi(b)$ or $\phi(-2b)=T^{\frac{q-1}{4}}(4)\phi(b)$. Note that $T^{\frac{q-1}{4}}(-1)=T^{\frac{q-1}{2}}(2)$ depends only on the congruence of $q \pmod 8$.
Thus, we have

$$N_{2,3}=q+qT^{\frac{q-1}{4}}\left(\frac{a^3}{27}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right).$$

Since $a_q(E_{2,3})=q-N_{2,3}$, we have

$$a_q(E_{2,3})=-qT^{\frac{q-1}{4}}\left(\frac{a^3}{27}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi\end{matrix} \Big| -\tfrac{27b^2}{4a^3}\right).$$

Example 4.2. *If $q\equiv 1 \pmod{36}$ and $E_{3,4}$ is an algebraic curve over $\mathbb{F}_q$, then the trace of Frobenius map on $E_{3,4}$ can be expressed as*

$$\begin{aligned}a_q(E_{3,4}(\mathbb{F}_q)) =& -T^{-\frac{q-1}{3}}(b)-T^{-\frac{2(q-1)}{3}}(b)-q^4\binom{T^{\frac{4(q-1)}{9}}}{T^{\frac{q-1}{3}}}\binom{T^{\frac{q-1}{36}}}{T^{\frac{5(q-1)}{36}}}T^{\frac{q-1}{3}}\left(\frac{3}{b}\right)\\ & \times {}_4F_3\left(\begin{matrix}\phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{5(q-1)}{9}}, & T^{\frac{2(q-1)}{9}}, & T^{\frac{8(q-1)}{9}}\end{matrix} \Big| -\tfrac{256b^3}{27a^4}\right)-q^4\binom{T^{\frac{5(q-1)}{9}}}{T^{\frac{2(q-1)}{3}}}\binom{T^{\frac{5(q-1)}{36}}}{T^{\frac{q-1}{36}}}\\ & \times T^{\frac{2(q-1)}{9}}(-1)T^{\frac{2(q-1)}{3}}\left(\frac{3}{b}\right){}_4F_3\left(\begin{matrix}\phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{4(q-1)}{9}}, & T^{\frac{q-1}{9}}, & T^{\frac{7(q-1)}{9}}\end{matrix} \Big| -\tfrac{256b^3}{27a^4}\right).\end{aligned}$$

Let $e=3$ and $d=4$ in Theorem 3.1. In this case we have,

$$\begin{aligned}N_{3,4}=& q+\sum_{i=1}^{2}T^{-\frac{i(q-1)}{3}}(b)+qT^{-\frac{7(q-1)}{12}}(-1)\sum_{i=1}^{2}M_iT^{\frac{i(q-1)}{3}}\left(\frac{3}{b}\right)\\ & \times {}_4F_3\left(\begin{matrix}\phi, & \epsilon, & \chi, & \chi^3 \\ & \psi^{6-i}, & \psi^{3-i}, & \psi^{9-i}\end{matrix} \Big| -\tfrac{256b^3}{27a^4}\right)\end{aligned} \tag{4.1}$$

where χ and ψ are characters of order 4 and 9 respectively, and

$$M_i = \left\{ G_{-\frac{i(q-1)}{3}} G_{-\frac{(6-i)(q-1)}{9}} (q-1) G_{\frac{(4i-3)(q-1)}{36}} G_{\frac{(4i-9)(q-1)}{36}} \right\}, i = 1, 2.$$

Using Lemmas 2.3 and 2.4, we find the values of M_i for $i = 1$,

$$M_1 = q^3 \binom{T^{\frac{4(q-1)}{9}}}{T^{\frac{q-1}{3}}} \binom{T^{\frac{q-1}{36}}}{T^{\frac{5(q-1)}{36}}} T^{\frac{7(q-1)}{12}}(-1),$$

and $i = 2$,

$$M_2 = q^3 \binom{T^{\frac{5(q-1)}{9}}}{T^{\frac{2(q-1)}{3}}} \binom{T^{\frac{5(q-1)}{36}}}{T^{\frac{q-1}{36}}} T^{\frac{29(q-1)}{36}}(-1).$$

By putting the value of M_i in equation (4.1), we have

$$\begin{aligned} N_{3,4} =& q + T^{-\frac{q-1}{3}}(b) + T^{-\frac{2(q-1)}{3}}(b) + q^4 \binom{T^{\frac{4(q-1)}{9}}}{T^{\frac{q-1}{3}}} \binom{T^{\frac{q-1}{36}}}{T^{\frac{5(q-1)}{36}}} T^{\frac{q-1}{3}} \left(\frac{3}{b}\right) \\ & \times {}_4F_3 \left(\begin{matrix} \phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{5(q-1)}{9}}, & T^{\frac{2(q-1)}{9}}, & T^{\frac{8(q-1)}{9}} \end{matrix} \Big| -\frac{256b^3}{27a^4} \right) + q^4 \binom{T^{\frac{5(q-1)}{9}}}{T^{\frac{2(q-1)}{3}}} \binom{T^{\frac{5(q-1)}{36}}}{T^{\frac{q-1}{36}}} \\ & \times T^{\frac{2(q-1)}{9}}(-1) T^{\frac{2(q-1)}{3}} \left(\frac{3}{b}\right) {}_4F_3 \left(\begin{matrix} \phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{4(q-1)}{9}}, & T^{\frac{q-1}{9}}, & T^{\frac{7(q-1)}{9}} \end{matrix} \Big| -\frac{256b^3}{27a^4} \right). \end{aligned}$$

Since $a_q(E_{3,4}) = q - N_{3,4}$, thus

$$\begin{aligned} a_q(E_{3,4}(\mathbb{F}_q)) =& - T^{-\frac{q-1}{3}}(b) - T^{-\frac{2(q-1)}{3}}(b) - q^4 \binom{T^{\frac{4(q-1)}{9}}}{T^{\frac{q-1}{3}}} \binom{T^{\frac{q-1}{36}}}{T^{\frac{5(q-1)}{36}}} T^{\frac{q-1}{3}} \left(\frac{3}{b}\right) \\ & \times {}_4F_3 \left(\begin{matrix} \phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{5(q-1)}{9}}, & T^{\frac{2(q-1)}{9}}, & T^{\frac{8(q-1)}{9}} \end{matrix} \Big| -\frac{256b^3}{27a^4} \right) - q^4 \binom{T^{\frac{5(q-1)}{9}}}{T^{\frac{2(q-1)}{3}}} \binom{T^{\frac{5(q-1)}{36}}}{T^{\frac{q-1}{36}}} \\ & \times T^{\frac{2(q-1)}{9}}(-1) T^{\frac{2(q-1)}{3}} \left(\frac{3}{b}\right) {}_4F_3 \left(\begin{matrix} \phi, & \epsilon, & T^{\frac{q-1}{4}}, & T^{\frac{3(q-1)}{4}} \\ & T^{\frac{4(q-1)}{9}}, & T^{\frac{q-1}{9}}, & T^{\frac{7(q-1)}{9}} \end{matrix} \Big| -\frac{256b^3}{27a^4} \right). \end{aligned}$$

We recall the following theorems from [3, 7].

Theorem 4.3. [3] *Let $q = p^n$, $p > 0$ an odd prime and let T be a generator of the character group $\widehat{\mathbb{F}_q^*}$. The number of points on the twisted Edward curve $C_{\alpha,\beta} : \alpha x^2 + y^2 = 1 + \beta x^2 y^2$ over $\mathbb{F}_q$ can be expressed as*

$$\#C_{\alpha,\beta}(\mathbb{F}_q) = q - 1 - \phi(\beta) - \phi(\alpha\beta) + q\phi(-\alpha) {}_2F_1 \left(\begin{matrix} \phi, & \phi \\ & \epsilon \end{matrix} \Big| \frac{\beta}{\alpha} \right).$$

Theorem 4.4. [7] *If A, B and C are characters of $\mathbb{F}_q$, then*

$${}_2F_1 \left(\begin{matrix} A, & \overline{A} \\ & \overline{A}B \end{matrix} \Big| \frac{1}{2} \right) = A(-2) \begin{cases} 0 & \text{if } B \text{ is not square} \\ \binom{C}{A} + \binom{\phi C}{A} & \text{if } B = C^2. \end{cases}$$

We have the following special case of the above theorem.

Corollary 4.5. *If $A = \phi$, $B = \phi$ and $q \equiv 1 (\mathrm{mod}\ 4)$, then we have*

$${}_2F_1 \left(\begin{matrix} \phi, & \phi \\ & \varepsilon \end{matrix} \Big| \frac{1}{2} \right) = \phi(-2) \left[\binom{T^{\frac{q-1}{4}}}{\phi} + \binom{T^{\frac{3(q-1)}{4}}}{\phi} \right].$$

Theorem 4.6. *Let $q = p^n$, $p > 0$ an odd prime and $q \equiv 1 (\mathrm{mod}\ 12)$. If $a, b \neq 0$, $k \in \mathbb{F}_q^*$ and $3k + a = 0$, then the number of points on the elliptic curve $E_{a,b,0} : y^2 = x^3 + ax^2 + bx$ can be expressed as*

$$\#E_{a,b,0} = q + qT^{\frac{3(q-1)}{4}} \left(\frac{3k^2 + 2ak + b}{3}\right) {}_2F_1 \left(\begin{matrix} T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi \end{matrix} \Big| -\tfrac{27(k^3+ak^2+bk)^2}{4(3k^2+2ak+b)^3} \right).$$

Proof. Since $a \neq 0$, we find $k \in \mathbb{F}_q^*$ such that $3k + a = 0$. A change of variables $(x, y) \mapsto (x + k, y)$ takes the algebraic curve $E_{a,b,0}$ to birationally equivalent form $E'_{2,3} : y^2 = x^3 + (3k^2 + 2ak + b)x + (k^3 + ak^2 + bk)$. Clearly $\#E_{a,b,0} = \#E'_{2,3}$, using Example 4.1 for $a' = 3k^2 + 2ak + b$ and $b' = k^3 + ak^2 + bk$, we have

$$\#E_{a,b,0} = q + qT^{\frac{3(q-1)}{4}} \left(\frac{3k^2 + 2ak + b}{3}\right) {}_2F_1 \left(\begin{matrix} T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi \end{matrix} \Big| -\tfrac{27(k^3+ak^2+bk)^2}{4(3k^2+2ak+b)^3} \right).$$

□

Corollary 4.7. *Let $q = p^n$, $p > 0$ an odd prime and $q \equiv 1 (\mathrm{mod}\ 12)$. If $k \in \mathbb{F}_q^*$ such that $3k + a = 0$ and $a \neq \pm 2\sqrt{b}$, then*

$$qT^{\frac{3(q-1)}{4}} \left(\frac{3k^2 + 2ak + b}{3}\right) {}_2F_1 \left(\begin{matrix} T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi \end{matrix} \Big| -\tfrac{27(k^3+ak^2+bk)^2}{4(3k^2+2ak+b)^3} \right) =$$
$$-\phi(a - 2\sqrt{b}) + \phi(ab - 2b\sqrt{b}) + q\phi(-a - 2\sqrt{b}) {}_2F_1 \left(\begin{matrix} \phi, & \phi \\ & \epsilon \end{matrix} \Big| \tfrac{a-2\sqrt{b}}{a+2\sqrt{b}} \right).$$

Proof. A change of variables $(x, y) \mapsto \left(\frac{x}{y}, \frac{x-\sqrt{b}}{x+\sqrt{b}}\right)$ takes the algebraic curve $E_{a,b,0} : y^2 = x^3 + ax^2 + bx$ to birationally equivalent form $C_{\alpha,\beta} : \alpha x^2 + y^2 = 1 + \beta x^2 y^2$, where $\alpha = a + 2\sqrt{b}$ and $\beta = a - 2\sqrt{b}$. Now the points on $E_{a,b,0}$ for $y = 0$ and $x = -\sqrt{b}$ do not correspond to any points on $C_{\alpha,\beta}$. For $y = 0$, there are $2 + T^{\frac{q-1}{2}}(a^2 - 4b)$ extra points on $E_{a,b,0}$, and for $x = -\sqrt{b}$, there are $1 + T^{\frac{q-1}{2}}(ab - 2b\sqrt{b})$ extra points on $E_{a,b,0}$. Similarly under the inverse transformation, a change of variables $(x, y) \mapsto \left(\frac{\sqrt{b}(1+y)}{1-y}, \frac{\sqrt{b}(1+y)}{x(1-y)}\right)$ takes the algebraic curve $C_{\alpha,\beta}$ to birationally equivalent form $E_{a,b,0}$. Now the points on $C_{\alpha,\beta}$ for $x = 0$ and $y = 1$ do not correspond to any points on $E_{a,b,0}$, so for $x = 0$ and $y = 1$, there are two extra points $(0, 1), (0, -1)$ on $C_{\alpha,\beta}$. Clearly $\#E_{a,b,0} + 2 = \#C_{\alpha,\beta} + 3 + T^{\frac{q-1}{2}}(a^2 - 4b) + T^{\frac{q-1}{2}}(ab - 2b\sqrt{b})$. From Theorems 4.3 and 4.6, we have

$$q + qT^{\frac{3(q-1)}{4}} \left(\frac{3k^2 + 2ak + b}{3}\right) {}_2F_1 \left(\begin{matrix} T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}} \\ & \phi \end{matrix} \Big| -\tfrac{27(k^3+ak^2+bk)^2}{4(3k^2+2ak+b)^3} \right) + 2 =$$
$$q - 1 - \phi(a - 2\sqrt{b}) - \phi(a^2 - 4b) + 3 + \phi(a^2 - 4b) + \phi(ab - 2b\sqrt{b})$$
$$+ q\phi(-a - 2\sqrt{b}) {}_2F_1 \left(\begin{matrix} \phi, & \phi \\ & \epsilon \end{matrix} \Big| \tfrac{a-2\sqrt{b}}{a+2\sqrt{b}} \right)$$

By solving the above equation, we have

$$qT^{\frac{3(q-1)}{4}}\left(\frac{3k^2+2ak+b}{3}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & \phi\end{matrix}\Big|-\tfrac{27(k^3+ak^2+bk)^2}{4(3k^2+2ak+b)^3}\right)=$$
$$-\phi(a-2\sqrt{b})+\phi(ab-2b\sqrt{b})+q\phi(-a-2\sqrt{b}){}_2F_1\left(\begin{matrix}\phi, & \phi\\ & \epsilon\end{matrix}\Big|\tfrac{a-2\sqrt{b}}{a+2\sqrt{b}}\right).$$

□

Corollary 4.8. *Let $q=p^n$, p be a prime with $q\equiv 1(\mathrm{mod}\ 12)$ and let T be a fixed generator of $\mathbb{F}_q^*$, then*

$${}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & \phi\end{matrix}\Big|\tfrac{1323}{1331}\right)=T^{\frac{q-1}{4}}\left(\frac{-44}{3}\right)\phi(2)\left[\binom{T^{\frac{q-1}{4}}}{\phi}+\binom{T^{\frac{3(q-1)}{4}}}{\phi}\right].$$

Proof. Set $a=12$ and $b=4$ in Corollary 4.7, we have

$$qT^{\frac{3(q-1)}{4}}\left(\frac{3k^2+24k+4}{3}\right){}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & \phi\end{matrix}\Big|-\tfrac{27(k^3+12k^2+4k)^2}{4(3k^2+24k+4)^3}\right)$$
$$=-\phi(8)+\phi(32)+q\phi(-16){}_2F_1\left(\begin{matrix}\phi, & \phi\\ & \epsilon\end{matrix}\Big|\tfrac{1}{2}\right).$$

Since in Corollary 4.7, $3k+a=0$, so k becomes -4, then we have

$${}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & \phi\end{matrix}\Big|\tfrac{1323}{1331}\right)=\phi(-1)T^{\frac{q-1}{4}}\left(\frac{-44}{3}\right){}_2F_1\left(\begin{matrix}\phi, & \phi\\ & \epsilon\end{matrix}\Big|\tfrac{1}{2}\right).$$

Using Corollary 4.5, the above equation becomes

$${}_2F_1\left(\begin{matrix}T^{\frac{q-1}{12}}, & T^{\frac{5(q-1)}{12}}\\ & \phi\end{matrix}\Big|\tfrac{1323}{1331}\right)=\phi(2)T^{\frac{q-1}{4}}\left(\frac{-44}{3}\right)\left[\binom{T^{\frac{q-1}{4}}}{\phi}+\binom{\frac{3(q-1)}{4}}{\phi}\right].$$

□

References

[1] Rupam Barman and Gautam Kalita. Certain values of gaussian hypergeometric series and a family of algebraic curves. *International Journal of Number Theory*, 8(04):945–961, 2012.

[2] Rupam Barman and Gautam Kalita. Hypergeometric functions and a family of algebraic curves. *The Ramanujan Journal*, 2(28):175–185, 2012.

[3] Rupam Barman and Gautam Kalita. Elliptic curves and special values of Gaussian hypergeometric series. *Journal of Number Theory*, 133(9):3099–3111, 2013.

[4] Rupam Barman and Gautam Kalita. On the polynomial x^d+ax+b over $\mathbb{F}_q$ and Gaussian hypergeometric series. *International Journal of Number Theory*, 09(07):1753–1763, 2013.

[5] Rupam Barman and Gautam Kalita. Hyperelliptic curves over $\mathbb{F}_q$ and Gaussian hypergeometric series. *Journal of the Ramanujan Mathematical Society*, 30(3):331–348, 2015.

[6] Jenny G Fuselier. *Hypergeometric functions over finite fields over $\mathbb{F}_p$ and relations to elliptic curves and modular forms*. PhD thesis, Texas A&M University, 2007.

[7] John Greene. Hypergeometric functions over finite fields. *Transactions of the American Mathematical Society*, 301(1):77–101, 1987.

[8] Masao Koike. Orthogonal matrices obtained from hypergeometric series over finite fields and elliptic curves over finite fields. *Hiroshima Mathematical Journal*, 25(1):43–52, 1995.

[9] Serge Lang. *Cyclotomic fields I and II*, volume 121. Springer Science and Business Media, 2012.

[10] Catherine Lennon. Gaussian hypergeometric evaluations of traces of Frobenius for elliptic curves. *Proceedings of the American Mathematical Society*, 139(6):1931–1938, 2011.

[11] Rudolf Lidl and Harald Niederreiter. *Finite fields*, volume 20. Cambridge university press, 1997.

[12] Ken Ono. Values of Gaussian hypergeometric series. *Transactions of the American Mathematical Society*, 350(3):1205–1223, 1998.

Department of Applied Mathematics, Indian Institute of Technology (ISM), Dhanbad-826004, Jharkhand, India

E-mail address: kewat.pk.am@ismdhanbad.ac.in, ramkumarbhu1991@gmail.com